

Linear Ordinary Differential Equation with Constant Coefficients

Amrik Sen

Engineering Mathematics in Action: FM112

10th January, 2022

Lecture Plan

Topic: Solving linear differential eqs. with constant coefficients

- 1 Conceptual introduction
- 2 Solution technique via examples
- 3 Ground work for harder problems

Form of Linear ODE w/ const. coeff.: $\mathcal{L}y(x) = 0$

$$\frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1 \frac{dy}{dx} + a_0 y = 0 \quad (1)$$

$$y^{(n)} + a_{n-1} y^{(n-1)} + \dots + a_1 y' + a_0 y = 0 \quad (2)$$

$\mathcal{L} := a_0 + a_1 \frac{d}{dx} + \dots + a_{n-1} \frac{d^{n-1}}{dx^{n-1}} + \frac{d^n}{dx^n}$ is the *linear differential operator*.

Solution form

Seek a solution of the form $y(x) = e^{rx}$ (Why?)

$\mathcal{L}(e^{rx}) = \mathcal{P}(r)e^{rx}$, where $\mathcal{P}(r) = r^n + a_{n-1}r^{n-1} + \dots + a_1 r + a_0$.

Characteristic equation

$$\mathcal{P}(r) = r^n + a_{n-1}r^{n-1} + \dots + a_1 r + a_0 = 0$$

Solution set depends on the nature of roots of characteristic eq.

- 1 n distinct real roots: $y(x) = c_1 e^{r_1 x} + c_2 e^{r_2 x} + \dots + c_n e^{r_n x}$
- 2 all real but some multiple roots: eg. $m \leq n$ multiple roots $r = r_0$, remaining roots are distinct; then
 $y(x) = (c_1 + c_2 x + \dots + c_m x^{m-1}) e^{r_0 x} + d_1 e^{r_1 x} + \dots + d_{n-m} e^{r_{(n-m)} x}$
 Thought exercise: justify why above is true?
 hint: $y = u(x) e^{r_0 x}$.
- 3 complex roots:
 $y = e^{\alpha x} (c_1 e^{i\beta x} + c_2 e^{-i\beta x}) +$ linear combination of real solutions

For repeated complex roots, follow the prescription in (2).

Example: ODEs with constant coeff. (distinct real roots)

Question: Solve the ODE $\frac{d^2y}{dt^2} + 5\frac{dy}{dt} + 6y = 0$ with initial conditions $y(0) = 1$, $\left. \frac{dy}{dt} \right|_{t=0} = 0$.

Example: ODEs with constant coeff. (distinct real roots)

Question: Solve the ODE $\frac{d^2y}{dt^2} + 5\frac{dy}{dt} + 6y = 0$ with initial conditions $y(0) = 1$, $\left. \frac{dy}{dt} \right|_{t=0} = 0$.

Solution: Characteristic equation:

$$r^2 + 5r + 6 = 0 \implies (r+2)(r+3) = 0 \text{ whence } r_{1,2} = -2, -3.$$

Example: ODEs with constant coeff. (distinct real roots)

Question: Solve the ODE $\frac{d^2y}{dt^2} + 5\frac{dy}{dt} + 6y = 0$ with initial conditions $y(0) = 1, \left. \frac{dy}{dt} \right|_{t=0} = 0$.

Solution: Characteristic equation:

$$r^2 + 5r + 6 = 0 \implies (r+2)(r+3) = 0 \text{ whence } r_{1,2} = -2, -3.$$

$$y(t) = c_1 e^{-2t} + c_2 e^{-3t}.$$

Example: ODEs with constant coeff. (distinct real roots)

Question: Solve the ODE $\frac{d^2y}{dt^2} + 5\frac{dy}{dt} + 6y = 0$ with initial conditions $y(0) = 1$, $\left. \frac{dy}{dt} \right|_{t=0} = 0$.

Solution: Characteristic equation:

$r^2 + 5r + 6 = 0 \implies (r+2)(r+3) = 0$ whence $r_{1,2} = -2, -3$.

$y(t) = c_1 e^{-2t} + c_2 e^{-3t}$. Now apply $y(0) = 1$ to obtain $c_1 + c_2 = 1$;
and $y'(0) = 0$ to obtain $-2c_1 - 3c_2 = 0$.

Example: ODEs with constant coeff. (distinct real roots)

Question: Solve the ODE $\frac{d^2y}{dt^2} + 5\frac{dy}{dt} + 6y = 0$ with initial conditions $y(0) = 1$, $\left. \frac{dy}{dt} \right|_{t=0} = 0$.

Solution: Characteristic equation:

$r^2 + 5r + 6 = 0 \implies (r+2)(r+3) = 0$ whence $r_{1,2} = -2, -3$.

$y(t) = c_1 e^{-2t} + c_2 e^{-3t}$. Now apply $y(0) = 1$ to obtain $\underline{c_1 + c_2 = 1}$; and $y'(0) = 0$ to obtain $\underline{-2c_1 - 3c_2 = 0}$. This gives $c_1 = 3$ and $c_2 = -2$.

Example: ODEs with constant coeff. (distinct real roots)

Question: Solve the ODE $\frac{d^2y}{dt^2} + 5\frac{dy}{dt} + 6y = 0$ with initial conditions $y(0) = 1$, $\left. \frac{dy}{dt} \right|_{t=0} = 0$.

Solution: Characteristic equation:

$r^2 + 5r + 6 = 0 \implies (r+2)(r+3) = 0$ whence $r_{1,2} = -2, -3$.

$y(t) = c_1 e^{-2t} + c_2 e^{-3t}$. Now apply $y(0) = 1$ to obtain $\underline{c_1 + c_2 = 1}$; and $y'(0) = 0$ to obtain $\underline{-2c_1 - 3c_2 = 0}$. This gives $c_1 = 3$ and $c_2 = -2$.

Thus $y(t) = 3e^{-2t} - 2e^{-3t}$.

Example: ODEs with constant coeff. (repeated real roots)

Question: Solve the ODE $y'' - 4y' + 4y = 0$ with initial conditions $y(0) = 1$, $y'(0) = 1$.

Example: ODEs with constant coeff. (repeated real roots)

Question: Solve the ODE $y'' - 4y' + 4y = 0$ with initial conditions $y(0) = 1$, $y'(0) = 1$.

Solution: Characteristic equation: $r^2 - 4r + 4 = 0 \implies (r - 2)^2 = 0$
i.e. double root: $r = 2$.

Example: ODEs with constant coeff. (repeated real roots)

Question: Solve the ODE $y'' - 4y' + 4y = 0$ with initial conditions $y(0) = 1$, $y'(0) = 1$.

Solution: Characteristic equation: $r^2 - 4r + 4 = 0 \implies (r - 2)^2 = 0$
i.e. double root: $r = 2$.

$$y(t) = c_1 e^{2t} + c_2 t e^{2t}.$$

Example: ODEs with constant coeff. (repeated real roots)

Question: Solve the ODE $y'' - 4y' + 4y = 0$ with initial conditions $y(0) = 1$, $y'(0) = 1$.

Solution: Characteristic equation: $r^2 - 4r + 4 = 0 \implies (r - 2)^2 = 0$
i.e. double root: $r = 2$.

$$y(t) = c_1 e^{2t} + c_2 t e^{2t}.$$

Now apply initial conditions to get: $c_1 = 1$, $c_2 = -1$.

Example: ODEs with constant coeff. (repeated real roots)

Question: Solve the ODE $y'' - 4y' + 4y = 0$ with initial conditions $y(0) = 1$, $y'(0) = 1$.

Solution: Characteristic equation: $r^2 - 4r + 4 = 0 \implies (r - 2)^2 = 0$
i.e. double root: $r = 2$.

$$y(t) = c_1 e^{2t} + c_2 t e^{2t}.$$

Now apply initial conditions to get: $c_1 = 1$, $c_2 = -1$.

Thus $y(t) = e^{2t} - t e^{2t}$.

Example: ODEs with constant coeff. (repeated real roots)

Question: Solve the ODE $y^{(v)} + 3y^{(iv)} + 3y''' + y'' = 0$ with some appropriate initial conditions (how many??).

Example: ODEs with constant coeff. (repeated real roots)

Question: Solve the ODE $y^{(v)} + 3y^{(iv)} + 3y''' + y'' = 0$ with some appropriate initial conditions (how many??).

Solution: Characteristic equation:

$r^5 + 3r^4 + 3r^3 + r^2 = (r+1)^3 r^2 = 0$ implies the roots are:
 $r = -1$ (triple root) and $r = 0$ (double root).

Example: ODEs with constant coeff. (repeated real roots)

Question: Solve the ODE $y^{(v)} + 3y^{(iv)} + 3y''' + y'' = 0$ with some appropriate initial conditions (how many??).

Solution: Characteristic equation:

$r^5 + 3r^4 + 3r^3 + r^2 = (r+1)^3 r^2 = 0$ implies the roots are:

$r = -1$ (triple root) and $r = 0$ (double root).

The solution will be of the form:

$$y(t) = (c_1 + c_2 t + c_3 t^2) e^{-t} + (c_4 + c_5 t) \dots$$

HW: complex roots

Question: Solve the ODE $\frac{d^4 y}{dt^4} + 8\frac{d^2 y}{dt^2} + 16y = 0$ with some appropriate initial conditions.

You may leave your answer in terms of constants of the problem.

Example: solving ODE with const. coeff.

Problem:

$$\varepsilon y'' + y = 0; \quad y(0) = 0, \quad y(1) = 1 \quad (3)$$

where, for now, ε is a constant.

Solution: Identify that the linear ODE has constant coefficients. Next, write the characteristic eq.:

$$r^2 + \frac{1}{\varepsilon} = 0 \quad (4)$$

Roots: $r = \pm \frac{i}{\sqrt{\varepsilon}}$. Therefore, $y(x) = c_1 e^{\frac{i}{\sqrt{\varepsilon}}x} + c_2 e^{-\frac{i}{\sqrt{\varepsilon}}x}$.

Then, apply boundary conditions:

$$y(0) = 0 \implies c_1 = -c_2 = c, \quad y(1) = 1 \implies c = \frac{1}{2\sin(1/\sqrt{\varepsilon})}.$$

Finally,
$$y(x) = \frac{\sin(x/\sqrt{\varepsilon})}{\sin(1/\sqrt{\varepsilon})}$$

Behavior of $y(x) = \frac{\sin(x/\sqrt{\varepsilon})}{\sin(1/\sqrt{\varepsilon})}$ as $\varepsilon \rightarrow 0^+$

Check: $\varepsilon = 1$ gives $y \sim \sin x$




Figure: $\varepsilon = 1.0$

Behavior of $y(x) = \frac{\sin(x/\sqrt{\varepsilon})}{\sin(1/\sqrt{\varepsilon})}$ as $\varepsilon \rightarrow 0^+$




Figure: $\varepsilon = 0.0001$

Behavior of $y(x) = \frac{\sin(x/\sqrt{\varepsilon})}{\sin(1/\sqrt{\varepsilon})}$ as $\varepsilon \rightarrow 0^+$




Figure: $\varepsilon = 0.0001$

For $\varepsilon = 0.0001$, we get rapid oscillations,
i.e. $y(x)$ exhibits discontinuity along x .

Singular Perturbation Problems: (a prelude to advanced mathematics for higher semesters)

Some options:

- 1 Since $\varepsilon \rightarrow 0^+$, ignore terms comprising ε ? Thus, the ODE $\varepsilon y'' + y = 0$ becomes $y = 0$. Clearly, $y(1) = 1$ contradicts $y(x) = 0$.

BAD OPTION!

- 2 WKB analysis: Seek solutions of the form

$$y(x) \sim e^{\frac{1}{\delta} \sum_{n=0}^{\infty} \delta^n S_n(x)}, \quad \delta \rightarrow 0 \quad (5)$$

Using (5) in $\varepsilon y'' + y = 0$, we obtain a hierarchy of closed differential equations for $S_n(x)$, solvable at every order of ε , to construct the asymptotic solution $y(x)$.